

Schizophrenia and the risk of fractures

Stubbs, Brendon; Gaughran, Fiona; Mitchell, Alex J.; De Hert, Marc; Farmer, Ross; Soundy, Andrew; Rosenbaum, Simon; Vancampfort, Davy

DOI:

[10.1016/j.genhosppsych.2015.01.004](https://doi.org/10.1016/j.genhosppsych.2015.01.004)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Stubbs, B, Gaughran, F, Mitchell, AJ, De Hert, M, Farmer, R, Soundy, A, Rosenbaum, S & Vancampfort, D 2015, 'Schizophrenia and the risk of fractures: a systematic review and comparative meta-analysis', *General Hospital Psychiatry*, vol. 37, no. 2, pp. 126-133. <https://doi.org/10.1016/j.genhosppsych.2015.01.004>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published as Stubbs Brendon, Gaughran Fiona, Mitchell Alex J., De Hert Marc, Farmer Ross, Soundy Andrew, Rosenbaum Simon, Vancampfort Davy, Schizophrenia and the risk of fractures: A systematic review and comparative meta-analysis, *General Hospital Psychiatry* (2015), doi: 10.1016/j.genhosppsych.2015.01.004

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Schizophrenia and the risk of fractures: A systematic review and comparative meta-analysis

Brendon Stubbs MSc, MCSP, Fiona Gaughran MD, MRCPsych, Alex J. Mitchell MD, MRCPsych, Marc De Hert MD, PhD, Ross Farmer BSc (Hons), MSc, Andrew Soundy PhD, Simon Rosenbaum PhD, Davy Vancampfort PhD, PT


PII: S0163-8343(15)00005-5
DOI: doi: [10.1016/j.genhosppsych.2015.01.004](https://doi.org/10.1016/j.genhosppsych.2015.01.004)
Reference: GHP 6959

To appear in: *General Hospital Psychiatry*

Received date: 3 November 2014
Revised date: 9 January 2015
Accepted date: 9 January 2015

Please cite this article as: Stubbs Brendon, Gaughran Fiona, Mitchell Alex J., De Hert Marc, Farmer Ross, Soundy Andrew, Rosenbaum Simon, Vancampfort Davy, Schizophrenia and the risk of fractures: A systematic review and comparative meta-analysis, *General Hospital Psychiatry* (2015), doi: [10.1016/j.genhosppsych.2015.01.004](https://doi.org/10.1016/j.genhosppsych.2015.01.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Schizophrenia and the risk of fractures: A systematic review and comparative meta-analysis

Brendon Stubbs¹, Fiona Gaughran², Alex J Mitchell³, Marc De Hert⁴, Ross Farmer⁵, Andrew Soundy⁶, Simon Rosenbaum^{7,8}, Davy Vancampfort^{4,9}

Submission to General Hospital Psychiatry

1. Faculty of Education and Health, University of Greenwich, Southwood Site Avery Hill Road
Eltham, London SE9 2UG, UK,
2. National Psychosis Service, South London and Maudsley NHS Foundation Trust, Denmark Hill,
London, UK
3. Department of Psycho-oncology, Leicester General Hospital, Leicestershire Partnership Trust,
Leicester, UK.
4. University Psychiatric Centre KU Leuven, Kortenberg, KU Leuven Departement of Neurosciences,
Leuvensesteenweg 517, B-3070 Kortenberg, Belgium
5. Physiotherapy department, South London and Maudsley NHS Foundation Trust, Denmark Hill,
London, UK
6. Department of Physiotherapy, University of Birmingham, Birmingham, B15 2TT, UK
7. Musculoskeletal Division, The George Institute for Global Health and School of Public Health,
University of Sydney, Sydney, Australia
8. School of Psychiatry, University of New South Wales, Sydney, Australia
9. KU Leuven Department of Rehabilitation Sciences, Tervuursevest 101, B-3001 Leuven, Belgium

Corresponding Author: Brendon Stubbs, University of Greenwich, New Eltham, Avery Hill Road,
London, SE9 2UG, United Kingdom. Telephone 004420 8331 8000, fax : 00441604696126 email
b.stubbs@greenwich.ac.uk

Abstract*Background*

People with schizophrenia experience increased rates of osteoporosis and may be at heightened risk of fractures. We conducted a systematic review and meta-analysis to investigate fractures among people with schizophrenia compared to people without mental illness.

Method

We systematically searched major electronic databases from inception till 10/2014. Articles were included that reported the number of fractures in people with schizophrenia and a control group. Two independent authors conducted searches, completed methodological assessment and extracted data. Data was narratively synthesised and a random effects incidence rate ratio (IRR) meta-analysis was performed.

Results

Eight studies were included encompassing 48,384 people with schizophrenia (49.9 -75.2 years, 41-100% female) and 3,945,783 controls. The pooled adjusted rate of fractures per 1000 person years was 5.54 (95% CI 4.92-5.57) in people with schizophrenia and 3.48 (95% CI 3.39-3.64) in control participants. The comparative meta-analysis showed that people with schizophrenia experience rate of fractures compared to control participants (IRR 1.72, 95% CI = 1.24 to 2.39, $I^2=49\%$; $n= 168,914$). There was insufficient data to conduct a robust moderator analysis, but the narrative review consistently highlighted antipsychotic medication was an important risk factor for fractures.

Conclusion

People with schizophrenia are at significantly increased risk of fractures. Future research is required to understand the mechanisms and should seek to validate fracture prediction algorithms used in the general population. Importantly there is a need to develop preventative strategies to improve

bone health and reduce fracture risk involving the wider multidisciplinary team and incorporating falls prevention strategies.

Key words: schizophrenia, psychosis, fracture, osteoporosis, fragility fracture

Introduction

Osteoporosis and associated fractures are a significant public health concern because of related mortality, morbidity, disability, and diminished quality of life ^{1,2}. Although osteoporosis is often referred to as the 'silent disease' it is highly clinically relevant due to the drastically increased risk of fractures which often occur as a consequence of the condition ^{1,2}. People with schizophrenia experience poorer physical health outcomes including reduced bone mineral density ³⁻⁶, with a recent systematic review establishing that people with schizophrenia are two and a half times more likely to have osteoporosis than people of similar age and sex without mental illness ⁷.

There are a multitude of complex reasons why people with schizophrenia may be at increased risk of fractures. For instance, people with schizophrenia typically take antipsychotic and other psychotropic medication which is associated with antipsychotic induced hyperprolactinaemia and osteoporosis ⁵ and falls ^{8,9} which are a leading cause of fracture due to trauma. Moreover, people with schizophrenia engage in lower levels of physical activity ¹⁰, have reduced lower limb strength ¹¹ and may experience high levels of pain ¹² which are important risk factors for falls ^{13,14} and therefore increase the risk of fracture. In addition, people with schizophrenia are at greatly increased risk of diabetes ^{15,16} which is a key risk factor for fractures in the general population ¹⁷. Some vulnerability for falling and fractures may derive from the illness itself since children who later go on to develop schizophrenia are noted to have more motor coordination difficulties than their peers ^{18,19}. Furthermore, alcohol use disorder is common in people with schizophrenia ⁶ and this may also increase risk of falling and subsequent fracture. Other potential risk factors that may also increase the risk of fractures include increased levels of stress hormones such as cortisol ²⁰⁻²².

It is particularly important to determine if there is a relationship between schizophrenia and fractures as research in general medicine has consistently established increased levels of mortality following fractures ²³. In addition, fractures in people with serious mental illnesses (SMI) such as schizophrenia can also lead to a deterioration of mental state ²⁴, higher post-operative infection

rates, worse ambulatory rates after one year, and a risk of contralateral fractures²⁵. Unfortunately, complications are more likely to arise in this group. A recent large study investigating 10,669,449 lower limb fractures²⁶ established that patients with schizophrenia (0.6% of the sample) spent more time in hospital post fracture (11 days) than any other patient group, including other groups such as dementia where there has been an increased effort to reduce fractures. In addition, Menendez et al²⁶ found that patients with schizophrenia experience significantly more adverse events following hip fractures including pneumonia, acute renal failure and deep venous thrombosis compared to those without mental illness (OR, 1.2; 95% CI, 1.2–1.3; $p < 0.001$).

Previously, a number of meta-analyses have reported that antipsychotic medication use is associated with an increased risk of fractures in older people^{8,27}. A selective narrative review in people with schizophrenia also indicated that osteoporotic fractures may have considerable adverse effects on general health, subjective well-being, the ability to engage in healthy lifestyle behaviors, and increased healthcare costs⁹. However, to date, no systematic review or meta-analysis has specifically investigated the relationship between schizophrenia and fractures. This is warranted in order to provide a rigorous up-to-date risk profile and inform relevant policy in this area. Therefore, we conducted a systematic review and meta-analysis to investigate the association between schizophrenia and fractures.

Method

This systematic review was conducted in accordance with the Meta-analysis of Observational Studies in Epidemiology guidelines²⁸ and reported in accordance with the PRISMA statement²⁹ following a predetermined but unpublished protocol.

Inclusion and exclusion criteria

We included observational studies (prospective, retrospective or cross-sectional) that: (a) included people with a diagnosis of schizophrenia according to recognised criteria (DSM V³⁰ or ICD 10³¹) together with a control group of people without a mental illness, and (b) reported the number of fractures over any period of time. We also included observational studies that reported the number of people with schizophrenia in comparative studies containing groups with and without a fracture. If we included mixed samples (e.g. pooled sample of SMI) we attempted to extract the schizophrenia specific data. If this was not possible we contacted the authors up to two times over a one month period to obtain the schizophrenia specific data. If we received no response and the study included $\geq 80\%$ with a diagnosis of schizophrenia we included the data. Due to the anticipated paucity of research, we collected data on any type and body site for the fracture that was confirmed through radiographs, medical note review or self-report. We did not place any language restrictions upon our searches. When we encountered studies reporting data from the same sample at different time points, we used the most recent data and/or the largest data set.

Information sources and searches

Two independent reviewers searched Academic Search Premier, MEDLINE, Psychology and Behavioral Sciences Collection, PsycINFO, SPORTDiscus, CINAHL Plus and Pubmed from inception until October 2014. We used the key words 'schizophrenia' or 'schiz*' or 'psychosis' and 'fracture*' or 'osteoporosis'. In addition, the reference lists of all eligible articles and recent systematic reviews

on bone health in people with schizophrenia were considered ^{7,9}. Primary/corresponding authors of research groups were contacted where necessary.

Study Selection

After the removal of duplicates, two independent reviewers screened the titles and abstracts of all potentially eligible articles. Both authors applied the eligibility criteria, and a list of full text articles was developed through consensus. The two reviewers then considered the full texts of these articles and the final list of included articles was reached through consensus.

Data Extraction

Two authors independently extracted data in a predetermined database. The data collected from each article included: study design, geographical location, details of schizophrenia participants (mean age, % males, diagnosis method, details of medications and chronicity of illness, and comparison group participant characteristics (mean age, % males)). We extracted data on fractures within the studies including the body site, method of acquiring the data, duration of data collection and details regarding the circumstances that contributed to fractures (e.g. falls, accidents).

Methodological quality assessment

Two authors completed methodological quality assessment of included articles using the Newcastle Ottawa Scale (NOS; ³²). The NOS is utilized to assess the methodological quality of non-randomized trials and has acceptable validity and reliability ³². The assessment tool focuses on three main methodological features: (1) the selection of the groups, (2) the comparability of the groups and (3) the ascertainment of the outcome of interest. Studies were given a NOS score ranging from 0-9, with a score of 5 or greater indicative of satisfactory methodological quality.

Data analysis

The results from the included studies were reported in a narrative synthesis and also a meta-analysis in accordance with the Cochrane reviewer's handbook ³³. Where possible we extracted raw data from the studies (or utilised data provided from authors upon request) regarding the number of people with and without a fracture in the schizophrenia and control groups. We then corrected for years of observation and sample size (i.e. person-years of observation) to compare the incidence rates (IR) of fractures across studies of differing time points and report this as a fracture rate per 1,000 years of observation per study. We pooled the data with a random effects meta-analysis calculating the incident rate ratio (IRR) to compare the rate of fracture between the two groups ³³. In accordance with the Cochrane reviewers handbook ³³ heterogeneity was assessed with the I^2 statistic and scores of 25%, 50% and 75% were classed as low, medium and high heterogeneity accordingly. We assessed publication bias with the visual inspection of a funnel plot ³³ and the begg ³⁴ and egger ³⁵ tests. Due to the anticipated dearth of studies and the heightened risk of obtaining a spurious result, we did not conduct moderator analysis with our results ³³. All analysis was conducted with Statsdirect and is presented with 95% confidence intervals (CI).

Results

Search results and study selection

The initial search identified 423 publications. After removal of duplicates, 367 abstracts and titles were screened (Figure 1). At the full text review stage, 44 articles were considered and 36 were subsequently excluded with reasons, leaving 8 articles that were included in the review. Details regarding the search results including reasons for exclusion of articles are summarised in figure 1.

Figure 1 here

Study and participant characteristics

Among the 8 included studies, 6 investigated the number of fractures among people with schizophrenia and a control group³⁶⁻⁴¹ including 48,384 people with schizophrenia and 3,945,783 people without mental illness. A further 2 studies reported the number of people with schizophrenia in a group of people with and without fractures (n=549 schizophrenia;^{42,43}). Of the included studies only 4 set out with the primary objective to investigate the relationship between schizophrenia and fractures^{36-38,42}. The summary of the groups of studies are presented in table 1 and 2 respectively. The average age among the participants ranged from 49.9 years³⁶ to 75.2 years³⁸. The fracture sites considered varied in each study but the hip was the most commonly investigated^{36-38,42,43}. None of the included studies contained clear information on the possible causes of fractures such as falls or accidents. Information regarding antipsychotic medication and other risk factors for osteoporosis and fractures was sparse among the included studies.

Table 1 and 2 here

Methodological quality

The NOS summary score for each article is presented in tables 1 and 2. All of the articles were of acceptable methodological quality with an average NOS score 6.25 (range 5-8).

Narrative results

Studies comparing the rate of fractures in people with schizophrenia and a control group

In a study among female patients with schizophrenia, Bishop et al⁴⁰ established that those with schizophrenia (13%) were more likely to have experienced a fracture in the past 12 months than the control group (0%, $p < 0.001$). More recently, Kelly et al⁴¹ investigated fractures over the past 12 months among 1,898 females with psychosis and 14,985 controls without any history of mental illness or substance use disorder. The authors⁴¹ established that similar proportions of people with psychosis (0.2%) and the control group had a fracture recorded in their medical records (0.3% control group). Jung et al³⁹ investigated the number of any fracture confirmed from radiograph among 229 inpatients with schizophrenia and established a higher rate among female patients ($N=24$, 25.8%) compared to the healthy controls of similar age (6.7%). The same results were evident among the male schizophrenia patients ($N=31$, 22.8%) compared to healthy male controls (4.6%).

Sorensen et al³⁶ conducted a national data linkage in Danish hospital registers investigating hip fractures including 15,431 patients with schizophrenia and 3,807,597 general population controls. The authors pooled data from 3,353,771 individuals and found that after adjusting for numerous established risk factors for fracture (sex, age, education, early retirement pension, physical comorbidity, lifetime corticosteroids, lifetime alcohol-related diagnosis, antidepressants, anticholinergics, and benzodiazepines) that a diagnosis with schizophrenia was associated with an increased risk for hip fracture ($IRR=1.19$, $95\%CI=1.08-1.31$). However, this effect was non-significant after the adjustment of antipsychotic medication ($IRR=1.00$, $95\%CI=0.90-1.11$). Sorensen et al³⁶ also investigated the impact of several medication classes consumed in the preceding 4 months on hip fracture and found in the unadjusted analysis that exposure to anticholinergics, antidepressants, benzodiazepines, antipsychotics (both low-potency, medium potency and high potency), olanzapine and quetiapine were all associated with a significantly increased risk of hip fracture. More specifically, when the authors compared the number of antipsychotic medications in the previous 4 months with those that had not taken any antipsychotic medication they found a dose response relationship of increased risk of hip fracture (IRR 1 antipsychotic= 1.30 ($95\%CI=1.10-1.54$), 2 antipsychotics 1.68 ($95\%CI=1.35-2.07$) and > 2 antipsychotics 1.81 ($95\%CI=1.28-2.54$). The authors

also demonstrated there was a dose response relationship of increased risk of hip fracture among those on long term prolactin (IRR=1.16, 95%CI=1.09–1.24) and non-prolactin raising antipsychotic medication (IRR=1.23, 95%CI=1.15–1.32).

In a large nationwide study, Tsai et al³⁷ investigated the number of major fractures among 30,335 people with schizophrenia and 121,340 age and gender matched controls across a ten year period. The authors established that 5.5% of patients with schizophrenia had a major fracture compared to 3.5% in the control group ($p<0.001$). The authors³⁷ established that patients with schizophrenia with higher psychiatric proportion of days covered (PDC= total number of medication covered days divided by the number of days in a certain time period) established that there was no dose response relationship, with an increasing PDC not indicating a higher fracture risk implying that the frequency of psychiatric medicine dose did not have an effect on major fracture risk. However, the authors were not able to disentangle the influence of specific groups of psychotropic medication on fracture risk. Across the 10 year study³⁷ period 473 people with schizophrenia (from 1673) died following a major fracture compared to 784 (from 4257) in the control group. The calculation of the RR from the raw data established that people with schizophrenia were at a 54% increased risk of mortality after a major fracture compared to the control group (RR=1.54, 95%CI=1.39 - 1.70). Lai et al³⁸ investigated hip fractures among a random sample of 445 patients with schizophrenia and 1,780 controls from the same cohort as Tsai et al³⁷. The authors found that people with schizophrenia were significantly more likely to experience hip fractures (IRR 1.91, 95% CI 1.49, 2.44).

Rate of fractures per 1000 person years

There was sufficient raw data from 5 studies to calculate the rate of fractures per 1000 person years³⁷⁻⁴¹. The rate of fractures per 1000 years were higher among participants with schizophrenia in 4 studies³⁷⁻⁴⁰ whilst they were slightly lower in Kelly et al⁴¹. Full details are presented in table 3. It was possible to pool the adjusted rate of fractures from 5 studies³⁷⁻⁴¹ in each group. This established that the rate of fractures was 5.54 (95% CI 4.92 to 5.57) fractures per 1000 person years

in those with schizophrenia and 3.48 (95% CI 3.39 to 3.64) fractures per 1000 person years in the control participants.

Table 3 here

Studies comparing the prevalence of schizophrenia in a fracture group and non-fracture group

Bolton et al⁴³ investigated hip, wrist and vertebral fractures from hospital records in Canada. The authors established that a diagnosis of schizophrenia is associated with fractures (OR=2.17, 95% CI 1.75-2.69). Howard et al⁴² investigated 16,341 hip fracture cases in a UK general database study against 29,889 non-hip fracture cases. The authors established that people with a diagnosis of schizophrenia were more likely to experience a hip fracture (OR=1.37, 95% CI 1.32-2.28). However, in the adjusted analysis when the authors investigated the influence of gender they found that male and female patients were not at increased risk of experiencing hip fractures (see table 3).

Meta-analysis

It was possible to pool data from five studies³⁷⁻⁴¹ involving 32,593 unique participants with schizophrenia and 138,186 control participants. This established an IRR of 1.72 (95% CI = 1.24 to 2.39, $I^2=49\%$; $n=168,914$) and is displayed in figure 2a. The funnel plot for the main analysis was broadly symmetrical (figure 2b) and the Begg-Mazumdar (Kendall's tau = 0.33 $P=0.46$) and Egger bias tests ($=0.49$ $P=0.44$) did not indicate any publication bias. In a subgroup analysis, we removed one study⁴¹ with very low fracture rates in both groups and found an IRR of = 1.82 (95% CI = 1.33 to 2.50, $I^2=47\%$, n schizophrenia=31,055, n controls=123,291).

Insert figure 2 and 2b here

Discussion

General findings

To our knowledge this is the first systematic review and meta-analysis to investigate fractures in people with schizophrenia. Across the 8 eligible studies the results are unequivocal indicating that people with schizophrenia are at increased risk of fractures. Both the results from the narrative review and meta-analyses suggest that the increased risk of fracture is approximately 50-100% in people with schizophrenia compared to people without mental illness. Clearly this is of concern, especially in light of the increased mortality rate among people with schizophrenia following major fractures³⁷ which may be attributed to the longer hospital stays and increased adverse events that people with schizophrenia experience following fractures²⁶.

The reasons for the increased risk of fractures observed in people with schizophrenia are likely to be complex and multifactorial. However, heightened levels of osteoporosis and reduced bone mass seen in this population⁷ are key factors since osteoporosis is a primary risk factor for fractures^{44,45}. Only one study³⁹ reported osteoporosis in both groups and they established a considerable increased risk in the patient group (34.9% v 18.4%). The results from the narrative review also implicate antipsychotic medication as being a key factor, which is not surprising given a previous meta-analysis has demonstrated an increased risk for older adults in receipt of antipsychotics⁸. In their large national database study, Sorensen et al³⁶ demonstrated a clear dose response relationship between the number of antipsychotics and fracture risk in the 4 months immediately prior to fracture. Interestingly, although antipsychotic hyperprolactinaemia has been proposed as one mechanism reducing bone mineral density and increasing fracture risk^{3,5,46}, Sorensen et al³⁶ established there was a comparable increased risk of fractures for those taking prolactin raising and sparing medication. However, it should be noted the effects of antipsychotics induced hyperprolactinaemia influence on bone mineral density will accumulate over many years^{5,47} and due to the age of participants it is likely that many participants in both groups may have taken 1st

generation antipsychotics in the past for several years and so bone mineral density may have already been influenced in those taking prolactin sparing medication. Other known risk factors for falls and fractures such as the presence of depression, polypharmacy, hypertension, diabetes and pain¹³ are all highly prevalent among people with schizophrenia yet, although sporadically reported in the current literature³⁷ there was insufficient data to clarify the importance of these factors on actual fractures.

Of the modifiable risk factors, lower limb muscle strength and balance are key predictors for falls and fractures in the general literature¹³ and these are known to be impaired in people with schizophrenia^{48,49}. Exercise is the most effective single intervention to prevent falls⁵⁰ and targeted exercise can improve lower limb strength, balance and may be integral in preventing falls and fractures in this patient group. When one considers that exercise is associated with a reduction in depressive symptoms and improved metabolic outcomes in people with schizophrenia⁵¹ and that exercise may be equally effective as pharmacological interventions to prevent cardiovascular disease related mortality⁵², then the central role of exercise in the treatment of schizophrenia is increasing. Research in the general medical literature⁵³ has demonstrated that better cardiorespiratory fitness is associated with shorter hospital stay and reduced mortality following major surgery. Recent research⁵⁴ has demonstrated that people with schizophrenia have greatly reduced cardiorespiratory fitness and exercise interventions should seek to improve this. To this end, physiotherapists and other specialists such as exercise physiologists may have particularly pertinent roles in the MDT management of schizophrenia⁵⁵. However, clearly future research is required to investigate and clarify this rationale. Interestingly, Bishop et al⁴⁰ established that people with schizophrenia were less likely than the control group to receive osteoporosis screening and pharmacological interventions. This is in line with a deficit in general preventative medical care seen in this group

^{56,57}.

Limitations

Despite the fact this is the first review of its kind, it is important to consider a number of limitations which are predominantly reflected by limitations in the primary data. First, due to the paucity of data and lack of information it was not possible to conduct moderator or subgroup analyses with the results. Specifically, we could not clearly investigate the influence of increasing age, different medication classes, duration of antipsychotic exposure and other important risk factors for fractures on the pooled results. However, this is partially appeased by the narrative results which included several very large age and sex matched studies adjusting for several confounders and still found an increased risk. Second, most of the studies were cross-sectional or retrospective cohort studies. Future prospective studies are required to further understand fractures in people with schizophrenia. Third, there was heterogeneity in the reporting of fractures and in the skeletal sites considered thus precluding evaluation of site-specific fracture risk across the studies.

Future research

There are many research questions of relevance to the clinical care of people with schizophrenia that need addressing as a priority. First, there is a need to better understand the reasons why people with schizophrenia are more likely to experience fractures and in particular investigate the contributing mechanisms, with a view to developing prevention plans. Also, research investigating the care of people with schizophrenia in general hospitals following fracture is warranted to investigate: (a) how they can be rehabilitated in the best manner and avoid prolonged stays in hospital and (b) why they appear more susceptible to adverse events in the hospital but also are at increased risk of mortality thereafter³⁷

In most of the studies we reviewed, the outcome was a fracture in any part of the skeleton and the circumstances in which the fracture incurred were not reported. Clearly, there is an urgent need to understand this in greater detail in order to prevent fractures in this group. There is a need for future prospective studies with specific outcome definitions, based on the site and the circumstances around the fracture to elucidate the potential mechanisms for the increased fracture

risk observed in people with schizophrenia. It is highly likely that heightened levels of osteoporosis and osteopenia reported in people with schizophrenia⁷ increase the risk for fractures. Investigations of mobility limitations and falls risk in people with schizophrenia should be considered since deficits in these areas are primary causes of falls and fractures in the general population^{44,45}. Future research could also consider the prospective investigation of falls to see if this is increased in people with schizophrenia compared to the general population.

Finally, our results show the importance of including fracture outcomes in prospective studies that provide individual patient level data in order to establish the attributable risk of schizophrenia on fracture, as well as the contribution of specific risk factors, including age, gender, race, weight, smoking, physical activity behavior (e.g. complying with health recommendations or not) and psychotropic medication use (e.g. use of long term prolactin raising medication or not, the use of benzodiazepines or not, etc.). Next to this, validation of fracture prediction algorithms used in the general population, such as FRAX⁵⁸ for people with schizophrenia might be an important area of future research to identify those most at risk of fracture.

Conclusion

Our systematic review and meta-analysis has demonstrated that people with schizophrenia are at increased risk of developing fractures. The exact reasons for this are not yet clear but antipsychotic medications are likely to contribute. Limitations in the primary data did not enable us to elucidate the underlying contributors and causes of fractures and there is a need for future prospective research to better understand this. In addition, in light of the concerns regarding increased morbidity and mortality among people with schizophrenia experiencing a fracture, there is a need to develop preventative interventions to stop fractures occurring and this should include the wider multidisciplinary team.

Conflict of Interest

BS, FG, AJM, RF, AS, SR have no conflict of interest related to this work.

Prof Dr De Hert has received consulting fees, speakers or advisory board fees, research support, or honoraria from AstraZeneca, Bristol- Myers Squibb, Eli Lilly, Janssen-Cilag, Lundbeck JA, Pfizer, and Sanofi- Aventis.

Dr Davy Vancampfort is funded by the Research Foundation - Flanders (FWO-Vlaanderen).

References

1. Kanis JA. Osteoporosis III: diagnosis of osteoporosis and assessment of fracture risk. *Lancet* 2002; **359**(9321): 1929-36.
2. Kanis J, McCloskey E, Johansson H, Cooper C, Rizzoli R, Reginster JY. European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporosis International* 2013; **24**(1): 23-57.
3. Hummer M, Malik P, Gasser RW, et al. Osteoporosis in patients with schizophrenia. *American Journal of Psychiatry* 2005; **162**(1): 162-7.
4. Leucht S, Burkard T, Henderson J, Maj M, Sartorius N. Physical illness and schizophrenia: a review of the literature. *Acta Psychiatrica Scandinavica* 2007; **116**(5): 317-33.
5. Stubbs B. Antipsychotic-induced hyperprolactinaemia in patients with schizophrenia: Considerations in relation to bone mineral density. *Journal of Psychiatric and Mental Health Nursing* 2009; **16**(9): 838-42.
6. De Hert M, Correll CU, Bobes J, et al. Physical illness in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in health care. *World Psychiatry: Official Journal Of The World Psychiatric Association (WPA)* 2011; **10**(1): 52-77.
7. Stubbs B, De Hert M, Sepehry AA, et al. A meta-analysis of prevalence estimates and moderators of low bone mass in people with schizophrenia. *Acta Psychiatrica Scandinavica* 2014.
8. Oderda LH, Asche CV, Pepper GA, Leipzig RM, Tasic D. Drugs and fall risk: an updated systematic review. *Communicating Nursing Research* 2009; **42**: 412-.
9. Kishimoto T, De Hert M, Carlson HE, Manu P, Correll CU. Osteoporosis and fracture risk in people with schizophrenia. *Current Opinion in Psychiatry* 2012; **25**(5): 415-29.
10. Soundy A, Wampers M, Probst M, et al. Physical activity and sedentary behaviour in outpatients with schizophrenia: A systematic review and meta-analysis... including commentary by Attux C, Leutwyler H, and Strohle A. *International Journal of Therapy & Rehabilitation* 2013; **20**(12): 588-96.
11. Vancampfort D, Probst M, Sweers K, Maurissen K, Knapen J, De Hert M. Relationships between obesity, functional exercise capacity, physical activity participation and physical self-perception in people with schizophrenia. *Acta Psychiatrica Scandinavica* 2011; **123**(6): 423-30.
12. Stubbs B, Mitchell, AJ., De Hert, M., Correll, C.U., Soundy, A., Stroobants, M., Vancampfort, D. The prevalence and moderators of clinical pain in people with schizophrenia: a systematic review and large scale meta-analysis. *Schizophrenia Research*; 2014 (in press).
13. Deandrea S, Lucenteforte E, Bravi F, Foschi R, La Vecchia C, Negri E. Risk factors for falls in community-dwelling older people: a systematic review and meta-analysis. *Epidemiology (Cambridge, Mass)* 2010; **21**(5): 658-68.
14. Stubbs B, Binnekade T, Eggermont L, Sepehry AA, Patchay S, Schofield P. Pain and the Risk for Falls in Community-Dwelling Older Adults: Systematic Review and Meta-Analysis. *Archives of Physical Medicine & Rehabilitation* 2014; **95**(1): 175-87.e9.
15. Vancampfort D, Wampers M, Mitchell AJ, et al. A meta-analysis of cardio-metabolic abnormalities in drug naïve, first-episode and multi-episode patients with schizophrenia versus general population controls. *World Psychiatry: Official Journal Of The World Psychiatric Association (WPA)* 2013; **12**(3): 240-50.
16. Mitchell AJ, Vancampfort D, De Herdt A, Yu W, De Hert M. Is the prevalence of metabolic syndrome and metabolic abnormalities increased in early schizophrenia? A comparative meta-analysis of first episode, untreated and treated patients. *Schizophrenia Bulletin* 2013; **39**(2): 295-305.
17. Janghorbani M, Van Dam RM, Willett WC, Hu FB. Systematic review of type 1 and type 2 diabetes mellitus and risk of fracture. *American Journal of Epidemiology* 2007; **166**(5): 495-505.
18. Dazzan P, Murray RM. Neurological soft signs in first-episode psychosis: A systematic review. *The British Journal of Psychiatry* 2002; **181**(Suppl43): s50-s7.

19. Rosso IM, Bearden CE, Hollister JM, et al. Childhood neuromotor dysfunction in schizophrenia patients and their unaffected siblings: a prospective cohort study. *Schizophrenia Bulletin* 2000; **26**(2): 367-78.
20. Abraham G, Halbreich U, Friedman RH, Josiassen RC. Bone mineral density and prolactin associations in patients with chronic schizophrenia. *Schizophrenia Research* 2003; **59**(1): 17-8.
21. Halbreich U, Rojansky N, Palter S, et al. Decreased bone mineral density in medicated psychiatric patients. *Psychosomatic Medicine* 1995; **57**(5): 485-91.
22. Halbreich U, Palter S. Accelerated osteoporosis in psychiatric patients: Possible pathophysiological processes. *Schizophrenia Bulletin* 1996; **22**(3): 447-54.
23. Sattui SE, Saag KG. Fracture mortality: associations with epidemiology and osteoporosis treatment. *Nature Reviews Endocrinology* 2014; **10**(10): 592-602.
24. Holt RIG. Osteoporosis in people with severe mental illness: A forgotten condition. *Maturitas* 2010; **67**(1): 1-2.
25. Javaid MK, Holt RIG. Understanding osteoporosis. *Journal of Psychopharmacology* 2008; **22**(2, Suppl): 38-45.
26. Menendez ME, Neuhaus V, Bot AGJ, Vrahas MS, Ring D. Do psychiatric comorbidities influence inpatient death, adverse events, and discharge after lower extremity fractures? *Clinical Orthopaedics And Related Research* 2013; **471**(10): 3336-48.
27. Hartikainen S, Lönnroos E. Use of sedatives and hypnotics, antidepressants and benzodiazepines in older people significantly increases their risk of falls. *Evidence Based Medicine* 2010; **15**(2): 59-.
28. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA: The Journal Of The American Medical Association* 2000; **283**(15): 2008-12.
29. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Clinical Trials* 2009; **6**(7): 1-6.
30. Association AP. Diagnostic and Statistical Manual of Mental Disorders – DSM-IV-TR. 4th edition. American Psychiatric Association; 2000.
31. organisation Wh. The ICD-10 Classification of Mental and Behavioural Disorders – Diagnostic Criteria for Research. 1993.
32. Wells G, Shea, B, O'Connell D, Peterson Jea. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses.
http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp (accessed 01/12/2013).
33. Higgins JPT, S G. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.
34. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994; **50**(4): 1088-101.
35. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ (Clinical Research Ed)* 1997; **315**(7109): 629-34.
36. Sørensen HJ, Jensen SOW, Nielsen J. Schizophrenia, antipsychotics and risk of hip fracture: A population-based analysis. *European Neuropsychopharmacology* 2013; **23**(8): 872-8.
37. Tsai K, Lee, CC., Chou, YM., Shen, SP., Su, CY., et al. The risks of major osteoporotic fractures in patients with schizophrenia: A population-based 10-year follow-up study. *Schizophrenia Research*; 2014 (in press).
38. Lai S-W, Lin C-L, Liao K-F. Higher incidence of hip fracture in newly diagnosed schizophrenic patients in Taiwan. *The Libyan Journal Of Medicine* 2013; **8**: 1-2.
39. Jung D-U, Kelly DL, Oh M-K, et al. Bone mineral density and osteoporosis risk in older patients with schizophrenia. *Journal of Clinical Psychopharmacology* 2011; **31**(4): 406-10.
40. Bishop JR, Alexander B, Lund BC, Klepser TB. Osteoporosis Screening and Treatment in Women with Schizophrenia: A Controlled Study. *Pharmacotherapy* 2004; **24**(4): 515-21.

41. Kelly DL, Myers CS, Abrams MT, et al. The impact of substance abuse on osteoporosis screening and risk of osteoporosis in women with psychotic disorders. *Osteoporosis International* 2011; **22**(4): 1133-43.
42. Howard L, Kirkwood G, Leese M. Risk of hip fracture in patients with a history of schizophrenia. *The British Journal of Psychiatry* 2007; **190**(2): 129-34.
43. Bolton JM, Metge C, Lix L, Prior H, Sareen J, Leslie WD. Fracture risk from psychotropic medications: A population-based analysis. *Journal of Clinical Psychopharmacology* 2008; **28**(4): 384-91.
44. Kanis JA. Diagnosis of osteoporosis and assessment of fracture risk. *Lancet* 2002; **359**(9321): 1929.
45. Kanis J, McCloskey E, Johansson H, Cooper C, Rizzoli R, Reginster Jy. European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporosis International* 2013; **24**(1): 23-57.
46. Takahashi T, Uchida H, John M, et al. The impact of prolactin-raising antipsychotics on bone mineral density in patients with schizophrenia: Findings from a longitudinal observational cohort. *Schizophrenia Research* 2013; **147**(2/3): 383-6.
47. Peuskens J, Pani L, Detraux J, De Hert M. The Effects of Novel and Newly Approved Antipsychotics on Serum Prolactin Levels: A Comprehensive Review. *CNS Drugs* 2014.
48. Vancampfort D, Probst M, De Hert A, et al. An impaired health related muscular fitness contributes to a reduced walking capacity in patients with schizophrenia: a cross-sectional study. *BMC Psychiatry* 2013; **13**: 5-.
49. Vancampfort D, Probst M, Scheewe T, et al. Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia. *Psychiatry Research* 2013; **207**(1-2): 25-32.
50. Sherrington C, Whitney JC, Lord SR, Herbert RD, Cumming RG, Close JCT. Effective exercise for the prevention of falls: a systematic review and meta-analysis. *Journal of the American Geriatrics Society* 2008; **56**(12): 2234-43.
51. Rosenbaum S, Tiedemann A, Sherrington C, Curtis J, Ward PB. Physical activity interventions for people with mental illness: a systematic review and meta-analysis. *The Journal Of Clinical Psychiatry* 2014.
52. Naci H, Ioannidis JPA. Comparative effectiveness of exercise and drug interventions on mortality outcomes: metaepidemiological study. *BMJ (Clinical Research Ed)* 2013; **347**: f5577-f.
53. Snowden CP, Prentis J, Jacques B, et al. Cardiorespiratory fitness predicts mortality and hospital length of stay after major elective surgery in older people. *Annals Of Surgery* 2013; **257**(6): 999-1004.
54. Vancampfort D, Guelinckx H, De Hert M, et al. Reliability and clinical correlates of the Astrand-Rhyming sub-maximal exercise test in patients with schizophrenia or schizoaffective disorder. *Psychiatry Research* 2014.
55. Stubbs B, Probst M, Soundy A, et al. Physiotherapists can help implement physical activity programmes in clinical practice. *The British Journal Of Psychiatry: The Journal Of Mental Science* 2014; **204**(2): 164-.
56. Lord O, Malone D, Mitchell AJ. Receipt of preventive medical care and medical screening for patients with mental illness: a comparative analysis. *General Hospital Psychiatry* 2010; **32**(5): 519-43.
57. **Mitchell A, I P, M Y, S P, V M, B S.** Is there a disparity in receipt of breast cancer screening among women with mental illness or distress? A comparative meta-analysis of mammography uptake. In press *British Journal of Psychiatry*; 2014.
58. Kanis JA, McCloskey EV, Johansson H, Oden A, Ström O, Borgström F. Development and use of FRAX in osteoporosis. *Osteoporosis International: A Journal Established As Result Of Cooperation Between The European Foundation For Osteoporosis And The National Osteoporosis Foundation Of The USA* 2010; **21 Suppl 2**: S407-S13.

Figure 1. PRISMA 2009 flow diagram for search strategy




Figure 2a Forest plot for the main meta-analysis




Figure 2b funnel plot for the main analysis




Table 1 – summary of included studies comparing the prevalence of fractures among people with schizophrenia compared to a control group

Study	Location and design	Schizophrenia Diagnosis	Schizophrenia Participants	Control participants	Fracture site and ascertainment	NOS score
Bishop et al 2004	US, retrospective study	DSMI IV	N=46 100% female 61.1 years	N=46 100% female 61.0 years	Fracture in past 12 months Medical records	5
Kelly et al 2011	US, retrospective cohort	ICD 9	N=1898 schizophrenia (psychosis) 100% female 56 years	N=14,985 healthy controls free from mental illness and substance use disorder 100% female 57 years	Any fracture in past 12 months Medical records	6
Jung et al 2011	Republic of Korea, cross sectional	DSM IV	N=229 inpatients 40.6% female Mean duration illness 142.6±79.2 months N=93 female, 59.1 years 91.4% FGA % 8.6% SGA N=136 male, 58.2 years 121 (89.0%) FGA and SGA 15 (11.0%) 34.9% osteoporosis	N=125 healthy controls 48% female N=60 female, 58.2 years N=65 males, 59.0 years 18.4% osteoporosis (p<0.01)	Any fracture across lifetime Medical records	7
Sorensen et al 2013	Denmark, Cohort study	ICD 8 and ICD 10	N= 15,431 41% female 49.9 years	N= 3,807,597 56.9% female 46.9 years	Hip fracture Medical records	8
Lai et al	Taiwan, cross	ICD 9	N=445	N= 1780	Hip fracture	5

2013	sectional		52.8% female 75.2 years	52.8% female 74.2 years	Medical records	
Tsai et al 2014	Taiwan, prospective cohort	ICD 9	N= 30,335 51.13% female 51.11 years	N= 121,340 51.13% female 51.11 years Age and gender matched no mental illness	Major fracture (vertebra, hip, humerus, forearm, wrist) Medical records	8

Key = FGA=first generation antipsychotic medication, SGA = second generation antipsychotic medication

Table 2 – Summary of included studies investigating the prevalence of schizophrenia in comparative studies of those with and without a fracture

Study	Location and design	Schizophrenia Diagnosis	Schizophrenia Participants with fracture	Schizophrenia participants without fracture	Fracture site and ascertainment	NOS score
<i>Howard et al 2007</i>	UK, retrospective cohort	GP records ICD 9	N=100 hip fracture cases with schizophrenia (0.61% of all cases)	N=110 people with schizophrenia non-hip fracture cases (0.37% of all non-fracture cases)	Hip fracture Medical records	6
<i>Bolton et al 2008</i>	Canada, retrospective cohort	ICD 9	N= 142 fracture cases with schizophrenia (0.9% of all fracture cases) No demographic data All > 50 years	N=197 people with schizophrenia did not have a fracture (0.4% of all non-fracture cases) All > 50 years	Hip, wrist vertebral Medical records	5

Table 3 – Results of included studies

Study	Type of fracture	Results
<i>Sorensen et al 2013</i>	Hip fracture	<p>Diagnosis of schizophrenia <u>IRR 1.19 (95% CI 1.08–1.31)^a</u></p> <p>Diagnosis of schizophrenia & adjusted for AP IRR 1.00 (95% CI 0.90–1.11)^a</p> <p>Medications taken in past 4 months prior to hip fracture</p> <p>Anticholinergics <u>IRR 1.46 (95% CI 1.26–1.70)^b</u></p> <p>Antidepressants <u>IRR 1.29 (95% CI 1.09–1.54)^b</u></p> <p>Benzodiazepines <u>IRR 1.34 (95% CI 1.14–1.58)^b</u></p> <p>Antipsychotics <u>IRR 1.42 (95% CI 1.21–1.65)^b</u></p> <p>Number of antipsychotics taken:</p> <p><u>1 IRR 1.30 (95% CI 1.10–1.54)^b</u></p> <p><u>2 IRR 1.68 (95% CI 1.35–2.07)^b</u></p> <p><u>>2 IRR 1.81 (95% CI 1.28–2.54)^b</u></p> <p>Long term prolactin raising antipsychotic medication (IRR 1.16 95% CI 1.09–1.24)</p> <p>Long term non prolactin raising antipsychotic medication (IRR 1.23 95% CI 1.15–1.32).</p>
<i>Bishop et al 2004</i>	Fracture in past 12 months	<p>Schizophrenia participants 130.43 (95% CI 47.86–283.90) fractures per 1000 person years</p> <p>Control participants 21.73 (95% CI 0.55–121.122) fractures per 1000 person years.</p> <p>IRR 6.00 (95% CI 0.72–275.98) calculated from raw data</p>
<i>Howard et al 2007</i>	Hip fracture	<p>Diagnosis of schizophrenia <u>OR 1.37 (95% CI 1.32–2.28)</u> unadjusted univariate</p> <p>Adjusted analysis:</p> <p>Female diagnosis with schizophrenia OR 1.01 (95% CI 0.72–1.40)</p> <p>Male diagnosis with schizophrenia OR 1.61 (95% CI 0.81–3.19)</p>
<i>Bolton et al 2008</i>	Hip, wrist, vertebral	<p>Diagnosis schizophrenia <u>OR 2.17 (95% CI 1.75–2.69)</u></p>
<i>Jung et al 2011</i>	Lifetime	Male participants with schizophrenia 3.86 (95% CI 2.62–5.48) fractures per 1000 person

	fracture	<p>years.</p> <p>Female participants with schizophrenia 4.10 (95% 2.62-6.10) fractures per 1000 person years</p> <p>Male control participants 0.782(95% CI 0.160-2.28) fractures per 1000 person years.</p> <p>Female control participants 1.12 (95% 0.30-2.89) fractures per 1000 person years.</p> <p>IRR 4.93 (95% 1.54-5.24) male participants</p> <p>IRR 3.63 (95% CI 1.24-14.40) female participants</p> <p>Calculated from raw data.</p>
<i>Kelly et al 2011</i>	Fracture in past 12 months	<p>Schizophrenia participants 2.10(95% CI 0.57-5.39) fractures per 1000 person years.</p> <p>Control participants 2.95 (95% 2.14 3.96) fractures per 1000 person years</p> <p>IRR calculated from raw data</p> <p>0.71 (95% 0.18- 1.95)</p>
<i>Lai et al 2013</i>	Hip fracture	<p>Schizophrenia participants 9.43 (95% CI 6.80-12.75) fractures per 1000 person years.</p> <p>Control participants 6.34 (95% CI 5.23-7.63) fractures per 1000 person years</p> <p>IRR for hip fracture among people with schizophrenia v control group</p> <p>IRR 1.91 (95% CI 1.49, 2.44)</p> <p>IRR calculated from raw data</p> <p><u>1.49 (95% CI 1.01-2.03)</u></p>
<i>Tsai et al 2014</i>	Major fracture Hip fracture	<p>Schizophrenia participants 5.52 (95% CI 5.26-5.79) major fractures per 1000 person years.</p> <p>Control participants 3.50 (95% CI 3.40-3.61) major fractures per 1000 person years</p> <p>Schizophrenia participants 2.66 (95% CI 2.47-2.85) hip fractures per 1000 person years.</p> <p>Control participants 1.02 (95% CI 0.96-1.07) hip fractures per 1000 years.</p> <p>IRR major fracture <u>1.57 (95% CI 1.48-1.66)</u> from raw data</p> <p>IRR hip fracture <u>2.60 (95% CI 2.38-2.85)</u> from raw data</p>

		Hazard ratio ^c for osteoporotic fracture investigating psychiatric proportion of days covered (PDC) an indicator of the intensity of drug exposure in patients with schizophrenia
	PDC = 0	HR 1.192 (95% CI 0.946–1.502)
	0 PDC ≤ 0.1	HR 1.190 (95% CI 0.968–1.464)
	0.1 PDC ≤ 0.2	<u>HR 1.373 (95% CI 1.059–1.782)</u>
	0.2 PDC ≤ 0.3	<u>HR 1.829 (95% CI 1.448–2.310)</u>
	0.3 PDC ≤ 0.4	<u>HR 1.829 (95% CI 1.448–2.310)</u>
	0.4 PDC ≤ 0.5	<u>HR 1.352 (95% CI 1.074–1.702)</u>
	0.5 PDC ≤ 0.6	<u>HR 1.695 (95% CI 1.388–2.071)</u>
	0.6 PDC ≤ 0.7	<u>HR 1.898 (95% CI 1.591–2.265)</u>
	0.7 PDC ≤ 0.8	<u>HR 1.563 (95% CI 1.298–1.883)</u>
	0.8 PDC ≤ 0.9	<u>HR 1.500 (95% CI 1.255–1.792)</u>
	0.9 b PDC ≤ 1	<u>HR 1.840 (95% CI 1.649–2.053)</u>

Key ^a (Sorensen et al 2013) adjusted for sex, age, education, early retirement pension, CPDS, lifetime corticosteroids, lifetime alcohol-related diagnosis, antidepressants, anticholinergics, and benzodiazepines ^b (Sorensen et al 2013) Adjusted for sex, age at diagnosis, alcohol misuse and somatic score. ^c (Tsai et al 2014) adjusted for age, gender, osteoporotic fracture-related illnesses, level of urbanization, and socioeconomic status